arXiv:1209.4844v1 [quant-ph] 21 Sep 2012

Parametric representation of open quantum systems and crossover from quantum to classical environment

D. Calvani,[1, 2] A. Cuccoli,[1, 2] N. Gidopoulos,[3] and P. Verrucchi[4, 1, 2]

[1] *Dipartimento di Fisica, Università di Firenze, Via G. Sansone 1, I-50019 Sesto Fiorentino (FI), Italy*
[2] *INFN Sezione di Firenze, via G.Sansone 1, I-50019 Sesto Fiorentino (FI), Italy*
[3] *ISIS, STFC, Rutherford Appleton Laboratory, Didcot, OX11 0QX, United Kingdom*
[4] *Istituto dei Sistemi Complessi, Consiglio Nazionale delle Ricerche,*
via G. Sansone 1, I-50019 Sesto Fiorentino (FI), Italy

(Dated: September 24, 2012)

We propose an approach to the study of open quantum systems based on a parametric representation of the principal system. The representation is obtained introducing generalized coherent states for the environment, and is such that the evolution from a fully quantum environment to a classical one can be inherently followed. The method is applied to the specific case where the principal system is a central qubit and the environment is a surrounding ring of interacting spins, with the overall Hamiltonian being that of the $s = 1/2$ Heisenberg star with frustration. In this case the quantum character of the environment, embodied by the total spin S of the ring, can be physically varied by acting on the ratio between the couplings entering the Hamiltonian, and formally followed by increasing S. We find that when the star is in any of its eigenstates, the qubit behaves as if it were under the effect of an external magnetic field, whose direction in real space is set by the variables defining the environmental coherent states, and is hence broadened according to their quantum probability distribution. When the quantum character of the environment is reduced, such distribution becomes narrower, finally collapsing into a Dirac-δ as $S \to \infty$. The entanglement between the central qubit and its environment remains well defined even in this limit, where it comes across as the binary entropy of the 2π-normalized Berry phase characterizing the dynamics of a qubit in an adiabatically precessing magnetic field.

Keywords: open quantum systems— central-spin models — Berry's phase

I. INTRODUCTION

Quantum systems with an environment are usually referred to as *open quantum systems* (OQS). Despite having been extensively studied since the very birth of quantum mechanics, the relevance acquired recently by certain features (such as coherence and entanglement) have boosted further the interest towards their behaviour [1, 2]. The description of an OQS in terms of the corresponding reduced density matrix, defined as the partial trace over the Hilbert space of the environment of the density matrix of the overall system, is axiomatically exact and fully retains the quantum character of the environment, making it the preferred approach, particularly in the realm of quantum information theory and computation [3, 4]. However, there exists another approach to OQS, where the principal system is in a pure state, but under the effect of a local Hamiltonian depending on "external" parameters, whose presence testifies the existence of a surrounding environment. At the heart of this approach, hereafter called "effective", stands the approximation that the environment be classical, so that the operators acting on its Hilbert space are replaced by c-number parameters; in this way, the interaction Hamiltonian is reduced to an effectively local one for the sole principal system. The prototypical example of this theoretical scheme is that of a spin in an external magnetic field, but many other examples can be easily spotted. Whenever dealing with one such situation, we will refer to the principal system as being "closed", according to the definition given in Ref. [1] and most often used in the literature. Notice that a closed system is not isolated, despite being described by a pure state.

In the exact and effective approaches sketched above, the environment is thus either fully quantum or completely classical. Aim of this work is that of devising a method for studying OQS such that an interpolation scheme becomes available, where the quantum character of the environment can be varied without affecting that of the principal system.

Being inspired by the representation which is typically used in quantum chemistry [5], where the electrons are described in terms of pure states parametrically dependent on the nuclei's coordinates, we devise a procedure for constructing a parametric representation for whatever OQS based on the use of generalized coherent states for its environment. This procedure is hence implemented in the case of a specific spin model, namely the spin-$\frac{1}{2}$ star with frustration[6, 7]: this system belongs to the large family of the so-called "central-spin" models, which have been extensively studied [8–20], since they describe magnetic interactions that play a relevant role in the physics of promising candidates for the realization of nanodevices [21–28].

The parametric representation, besides providing an original insight into the physical behaviour of the above spin model, prove suitable for dealing with peculiar geometrical effects such as the Berry's phase[29, 30], and allows us to confront a very relevant topic pertaining OQS, namely the intimate connection between geometric aspects of quantum mechanics and entanglement[31, 32], as thoroughly discussed in the final part of the paper.

II. THE PARAMETRIC REPRESENTATION WITH COHERENT STATES

Let us consider an isolated system in a pure, normalized, state $|\Psi\rangle$: the system is made of a principal system A and its environment B, with Hilbert space $\mathcal{H}_A$ and $\mathcal{H}_B$, respectively, both of finite dimension. Given two orthonormal bases $\{|\alpha\rangle\} \subset \mathcal{H}_A$ and $\{|\beta\rangle\} \subset \mathcal{H}_B$, it is $|\Psi\rangle = \sum_{\alpha\beta} c_{\alpha\beta}|\alpha\rangle|\beta\rangle$. Aimed at devising the interpolation scheme mentioned in the Introduction, we understand that describing the environment in terms of continuous parameters, rather than with a set of discrete ones, can be rewarding. The idea of introducing proper "environmental coherent states" naturally follows, and we accomplish it by resorting to the group-theoretical construction proposed by Gilmore (see Ref.[33] and references therein). Let the total environmental Hamiltonian be $H_B + H_{AB}$, where the first, local, term contains operators acting on $\mathcal{H}_B$, while the second, interaction, term contains operators acting on $\mathcal{H}_A \otimes \mathcal{H}_B$. We assume H_{AB} is a linear combination of tensor products of operators acting on $\mathcal{H}_A$ and on $\mathcal{H}_B$, as is the case in most physical situations. Let G be the dynamical group in terms of whose generators we can write the environmental Hamiltonian, and $\mathfrak{g}$ be its algebra; the Hilbert space $\mathcal{H}_B$ is given by the specific physical set up and is associated with a unitary irreducible representation of G. The choice of a reference state $|\beta_0\rangle \in \mathcal{H}_B$ fixes: *i)* the maximum stability subgroup $F \subset G$, i.e. the set of those operators f such that $f|\beta_0\rangle = e^{i\lambda_f}|\beta_0\rangle$; *ii)* the quotient G/F, such that any $g \in G$ can be locally decomposed as $g = \Omega f$, with $f \in F$ and $\Omega \in G/F$. Generalized *coherent states* $|\Omega\rangle$ are eventually defined as $|\Omega\rangle = \Omega|\beta_0\rangle$, with $\Omega \in G/F$; they can be normalized, and form an overcomplete set in $\mathcal{H}_B$, *i.e.*

$$\int \mathrm{d}\mu(\Omega)|\Omega\rangle\langle\Omega| = \mathbf{1}_{\mathcal{H}_B} \ , \tag{1}$$

with $\mathrm{d}\mu(\Omega)$ a proper group-invariant measure in G/F, and $\langle\Omega|\Omega'\rangle \neq \delta(\Omega - \Omega')$. Using the above resolution of the identity in $\mathcal{H}_B$, the state of the total system can be written as

$$|\Psi\rangle = \int \mathrm{d}\mu(\Omega)\chi^\Psi(\Omega)|\Omega\rangle|\phi_A^\Psi(\Omega)\rangle \ , \tag{2}$$

with

$$|\phi_A^\Psi(\Omega)\rangle \equiv \frac{1}{\chi^\Psi(\Omega)}\sum_\alpha f_\alpha(\Omega)|\alpha\rangle \ , \tag{3}$$

$$f_\alpha(\Omega) \equiv \sum_\beta \langle\Omega|\beta\rangle c_{\alpha\beta} \ , \tag{4}$$

$$\chi^\Psi(\Omega) \equiv e^{i\Lambda(\Omega)}\sqrt{\sum_\alpha |f_\alpha(\Omega)|^2} \ , \tag{5}$$

where $e^{i\Lambda(\Omega)}$ represents the gauge freedom pertaining the definition of the coherent state $|\Omega\rangle$. Each ket $|\phi_A^\Psi(\Omega)\rangle$ is a normalized element of $\mathcal{H}_A$ and therefore represents a physical state for the principal system. By parametric representation of A we will hereafter mean its description in terms of the pure states $\{|\phi_A(\Omega)\rangle\}$: the dependence of $|\phi_A(\Omega)\rangle$ on the parameter Ω is the fingerprint that an environment exists. Actually, it can be easily shown that $|\phi_A^\Psi\rangle$ depends on the parameter Ω if and only if the global state $|\Psi\rangle$ is entangled: it is the entangled structure of $|\Psi\rangle$ that causes the dependence on Ω to be conveyed from the environment to the principal system, thus resulting as the ultimate responsible for the emergence of phenomena related with the geometrical properties of the parameters space, as shown below in the case of the Berry's phase. The normalization of $|\Psi\rangle$ implies $\int \mathrm{d}\mu(\Omega)|\chi^\Psi(\Omega)|^2 = 1$, allowing $|\chi^\Psi(\Omega)|^2$ to be interpreted as a probability distribution on G/F, which turns out to be the phase space of the environment under rather general assumptions[33]; indeed, by a simple calculation it is immediate to show that $|\chi^\Psi(\Omega)|^2$ is just the Husimi Q-function of the environment reduced density matrix [33, 34], $\langle\Omega|(\mathrm{Tr}_A\,|\Psi\rangle\langle\Psi|)|\Omega\rangle = |\chi^\Psi(\Omega)|^2$.

The resulting picture for the principal system is that of a continuouos collection of pure, parametrized, states whose occurrence is ruled by a probability distribution over the phase space of the environment. The relation $\mathrm{Tr}_B[\,\cdot\,] = \int \mathrm{d}\mu(\Omega)\langle\Omega|\,\cdot\,|\Omega\rangle$ holds [34], yielding, for the reduced density matrix of the principal system, $\rho_A = \int \mathrm{d}\mu(\Omega)|\chi(\Omega)|^2|\phi(\Omega)\rangle\langle\phi(\Omega)|$, and hence

$$\langle O\rangle \equiv \mathrm{Tr}_A(\rho_A O) = \int \mathrm{d}\mu(\Omega)|\chi(\Omega)|^2\langle\phi(\Omega)|O|\phi(\Omega)\rangle \ , \tag{6}$$

for any principal-system observable O. The above construction allows us to treat the principal system and its environment in a very different formal scheme: in fact, and at variance with other related works[11, 35], coherent states are here exclusively adopted for the environment. This makes the approximations which naturally arise in the coherent state formalism available for describing the environment, but it also prevents the principal system from being affected by those same approximations, which is, as shown below for the classical limit, an essential feature of our approach.

Before ending this section, we notice that any resolution of the identity in the Hilbert space of the environment defines a parametric representation for the principal system. However, if the resolution implies summing over a set of discrete indexes, the representation coincides with that obtained by the Schmidt decomposition. It is indeed the possibility of using continuous parameters (which implies introducing a proper quantum phase space for the environment) that gives the above construction a surplus value. To this respect, the construction of the parametric representation described above can be both generalized, referring to approaches to quantum mechanics on phase space [36] that go beyond the theory of coherent states, and made more specific, whenever continuous variables different from those related to coherent states emerge in dealing with specific types of environment [37, 38].

III. THE SPIN-$\frac{1}{2}$ STAR WITH FRUSTRATION.

We apply the above formalism to the specific physical situation where a spin 1/2 ($\boldsymbol{\sigma}/2$, hereafter called *qubit*) interacts with an even set of N spins 1/2 ($\boldsymbol{s}_i$ hereafter called *environmental spins*) *via* an isotropic antiferromagnetic coupling [6, 7]. The environmental spins interact among themselves and the total Hamiltonian is that of the so-called "spin-$\frac{1}{2}$ star with frustration",

$$H = H_B + H_{AB} \ , \tag{7}$$

$$H_B = \frac{k}{2N}\sum_i^N \boldsymbol{s}_i \cdot \boldsymbol{s}_{i+1} \ \ ; \quad k > 0 \tag{8}$$

$$H_{AB} = \frac{g}{2N}\frac{\boldsymbol{\sigma}}{2}\cdot\sum_i \boldsymbol{s}_i \ \ ; \quad g > 0 \tag{9}$$

where i runs over the sites of the external ring. As mentioned in the introduction, the spin-$\frac{1}{2}$ star with frustration belongs to a class of "central-spin"-like models that have been extensively studied in the last decade. Without entering a detailed case study, which goes beyond the scope of this paper, we underline that the spin-$\frac{1}{2}$ star with frustration posseses the most welcome property of allowing the quantum character of the environment, as measured by the total spin of the ring, to be varied acting on the ratio k/g (usually referred to as the *frustration ratio*), as discussed below. Moreover the model is exactly solvable and analytical expressions of its eigenstates and eigenvalues are available [6].

Let us here briefly review the main known facts about the spin-$\frac{1}{2}$ star with frustration, Hamiltonian (7)-(9): the integrals of motion are the total Hamiltonian H, the local environmental Hamiltonian H_B, the square $\boldsymbol{J}^2$ of the total angular momentum $\boldsymbol{J} \equiv \boldsymbol{\sigma}/2 + \sum_i \boldsymbol{s}_i$, its component along the quantization axis, J^z, and the square $\boldsymbol{S}^2$ of the total spin of the environment $\boldsymbol{S} = \sum_i \boldsymbol{s}_i$, with the respective eigenvalues E, kE_B, $J(J+1)$, M, and $S(S+1)$; it is also useful to define m as the eigenvalues of the component S^z of the total environmental spin $\boldsymbol{S}$. The relations $J = S \pm \frac{1}{2}$ and $M = m \pm \frac{1}{2}$ hold. The total spin S ranges from 0 to $N/2$. Apart from the state with $S = 0$, which has energy kE_B, for each assigned value $S > 0$ the energy spectrum consists of the two multiplets

$$E_+ \ = \ kE_B + \frac{g}{2N}S \ , \tag{10}$$

$$E_- \ = \ kE_B - \frac{g}{2N}(S+1) \ ; \tag{11}$$

the lowest eigenvalue of H_B obeys the Lieb-Mattis ordering [39], $E_B(S) < E_B(S+1)$, and the competition between the two terms in Eq. (11), embodying the antiferromagnetic frustration of the model, makes the groundstate (GS) of the star belong to the E_- multiplet with a value of S that varies with the frustration ratio k/g. In fact, for $0 \leq k/g \leq 1/4 \equiv \alpha_0$, the GS has $S = N/2$ while, when k/g increases, there exists a sequence of critical values α_n, $n = 1, ..., (N/2-1)$, such that $S = N/2 - n$ for $\alpha_{n-1} < k/g \leq \alpha_n$, and $S = 0$ for $k/g > \alpha_{N/2-1} >> 1$. Notice that the critical values α_n depend on N, with the exception of α_0 which equals 1/4 for all N. The basic structure of the eigenstates belonging to a subspace with fixed S is

$$|\Psi_M^\pm\rangle = a_M^\pm|\uparrow\rangle|m_-\rangle + b_M^\pm|\downarrow\rangle|m_+\rangle \ , \tag{12}$$

where $|\uparrow,\downarrow\rangle$ are the eigenstates of σ^z, $|m_\pm\rangle$ are the eigenstates of S^z with eigenvalues $m = M \pm 1/2$, and the apex $\pm$ refers to the state having energy $E_\pm$. Introducing $\widetilde{S} \equiv S + 1/2$, the coefficients are

$$a_M^\pm = \pm\frac{1}{\sqrt{2}}\sqrt{1 \pm \frac{M}{\widetilde{S}}} \ , \ b_M^\pm = \frac{1}{\sqrt{2}}\sqrt{1 \mp \frac{M}{\widetilde{S}}} \ , \tag{13}$$

yielding $a^\pm = \pm b^\mp$. The entanglement between the qubit and its environment for the states (12), as measured by the Von Neumann entropy, is easily found to be the same for both multiplets and to depend just on the ratio $M/\widetilde{S}$, according to

$$\mathcal{E}_{\sigma S} = -h\left[\frac{1}{2}\left(1 - \frac{M}{\widetilde{S}}\right)\right] \ , \tag{14}$$

where $h[x]$ $(0 \leq x \leq 1)$ is the binary entropy $x\log x + (1-x)\log(1-x)$.

We construct the parametric representation of the states (12), each having S fixed, by identifying $\boldsymbol{S} = \sum_i \boldsymbol{s}_i$ with the environment and $\boldsymbol{\sigma}$ with the principal system; the dynamical group G to be employed in the generalized coherent state construction is therefore $SU(2)$, with $\mathcal{H}_B$ its spin$-S$ irreducible representation. We choose the reference state $|\boldsymbol{\beta}_0\rangle$ to be the maximal weight of the spin S-representation, *i.e.* $|\boldsymbol{\beta}_0\rangle = |m = S\rangle \in \mathcal{H}_B$, so that $U(1)$ is the maximum stability subgroup F, eventually ending up in the *Bloch coherent states* $|\Omega\rangle$, whose expansion over the S_z basis $\{|m\rangle\} \in \mathcal{H}_B$ is (using the conventions of [34])

$$|\Omega\rangle \equiv e^{\zeta S^- - \zeta^* S^+}|m = S\rangle = \sum_{m=-S}^{S} g_m(\theta)e^{\mathrm{i}(S-m)\varphi}|m\rangle \ , \tag{15}$$

where $S^\pm = S_x \pm iS_y$, $\zeta = \frac{\theta}{2}e^{i\varphi}$,

$$g_m(\theta) \equiv \sqrt{\binom{2S}{S+m}}\cos^{S+m}\left(\frac{\theta}{2}\right)\sin^{S-m}\left(\frac{\theta}{2}\right) \ , \tag{16}$$

and $\Omega \equiv (\theta, \varphi)$ is a point of the $S^2 \simeq SU(2)/U(1)$ sphere. The resolution of the identity reads $\mathbf{1}_{\mathcal{H}_B} = (\widetilde{S}/2\pi)\int \mathrm{d}\Omega|\Omega\rangle\langle\Omega|$, where $\mathrm{d}\Omega \equiv \sin\theta \mathrm{d}\theta\mathrm{d}\varphi$ is the euclidean measure on the S^2 sphere. From the definitions

(2)-(5) and the expansion (15), we can explicitely write the states (12) as

$$|\Psi_M^\pm\rangle = \frac{\widetilde{S}}{2\pi}\int \mathrm{d}\Omega\, \chi_M^\pm(\Omega)|\Omega\rangle|\phi_M^\pm(\Omega)\rangle\ , \tag{17}$$

where

$$\chi_M^\pm(\Omega) = e^{\mathrm{i}(\widetilde{S}-M)\varphi}\sqrt{[a_M^\pm g_{M-\frac{1}{2}}(\theta)]^2 + [b_M^\pm g_{M+\frac{1}{2}}(\theta)]^2}\ , \tag{18}$$

and the qubit parametrized states are

$$|\phi_M^\pm(\Omega)\rangle = \cos\left(\frac{\Theta_M^\pm(\theta)}{2}\right)|\uparrow\rangle + \sin\left(\frac{\Theta_M^\pm(\theta)}{2}\right)e^{\mathrm{i}\varphi}|\downarrow\rangle\ , \tag{19}$$

where $\Theta_M^\pm(\theta)$ are the solution with respecto to Θ of the equations

$$\tan\frac{\Theta}{2} = \left(\tan\frac{\theta_M}{2}\right)^{\pm 1}\tan\frac{\theta_M}{2}\cot\frac{\theta}{2}\ , \tag{20}$$

and $\theta_M \equiv \arccos(M/\widetilde{S})$. The parametric representation of the qubit is provided by the kets (19), each corresponding to the physical state the qubit is in, when the total system is in $|\Psi_M^\pm\rangle$ and the environment is the coherent state $|\Omega\rangle$. As $|\chi_M^\pm(\Omega)|$ does not depend on φ, the Ω-occurence is ruled by the θ-normalized probability distribution $\widetilde{S}\sin(\theta)|\chi_M^\pm(\Omega)|^2 \equiv p_M^\pm(\theta)$. In the upper panel of Fig.1 we show $p_M^-(\theta)$ for different values of $M/\widetilde{S}$; the values of S are chosen so as to consistently correspond to half-integer M. Two effects clearly testifies that we are dealing with a quantum environment: the finite width of the distribution and the position shift of its maximum with respect to θ_M. When the quantum character of the environment is reduced, *i.e.* S is increased with $M/\widetilde{S}$ fixed, both quantities lessen. Notice that the above shift, and consequently the shape of $p_M^-(\theta)$, markedly depend on $M/\tilde{S}$, due the distribution being defined, and normalized, on the compact support $[0,\pi]$.

The probability distribution of the qubit states (19) on the Bloch sphere, parametrized by Θ and φ, is the integrand of Eq.(6) with $O = |\phi_M^\pm\rangle\langle\phi_M^\pm|$, and reads $\pi_M^\pm(\Theta,\varphi) = p_M^\pm(\theta)$, with Θ and θ connected by Eq.(20). When the total system is in its GS multiplet, it is easily seen that $\Theta = \pi - \theta$ for all values of M. The lower panel of Fig.1 shows $\pi_M^-(\Theta,\varphi) = p_M^-(\pi-\Theta)$ for $M/\widetilde{S}$ as in the upper panel, and $S = 5$ (each sphere is below the corresponding distribution); the angle θ_M is identified by the black line of latitude. The conditional probability distribution for the qubit to be, say, in the state $|\uparrow\rangle$, reads

$$y_M^{\pm,\uparrow}(\theta) = p_M^\pm(\theta)\cos^2\frac{\Theta_M^\pm(\theta)}{2}\ ; \tag{21}$$

In the case of the GS multiplet, despite $|\phi_M^-(\Omega)\rangle$ being independent on M, due to $\Theta_M^- = \pi - \theta$ for all M, the conditional probability distribution (21) inherits the dependence on M from the environment, as cleary seen in Fig.2. Notice that the dependence on M accounts for the antiferromagnetic character of the interaction term (9) by representing the counteralignment of the qubit with respect to the total environmental spin. Let us now further comment on the parametrized qubit states (19): the kets $|\phi_M^\pm(\Omega)\rangle$ are apparently identified as ground ("-")



(a)$p_M^-(\theta)$ for varying S and θ_M



(b)$M = 9/2$ (c)$M = 3/2$ (d)$M = -3/2$ (e)$M = -9/2$

FIG. 1: Upper panel: $p_M^-(\theta)$ for $\frac{M}{\widetilde{S}} = \frac{9}{11}, \frac{3}{11}, -\frac{3}{11}, -\frac{9}{11}$ (from left to right) and S=5,16,27,38,49 (from below); the vertical dashed lines mark the corresponding values of θ_M (see text), each identified by a given color. Lower panel: qubit-states distributions, $\pi_M^-(\pi-\theta,\varphi)$, on the Bloch sphere for $S = 5$ and θ_M as in the upper panel. A black line of latitude marks θ_M on each sphere, and the corresponding value of M is reported below.



FIG. 2: Conditional probability distributions for the qubit to be in the $|\uparrow\rangle$ state when the total system is in its ground state. Values of S and $M/\widetilde{S}$, as well as dashed lines, as in Fig. 1.

and excited ("+") states of proper Zeeman terms, whose explicit form can be obtained by requiring that their corresponding energy, according to Eq.(6), be equal to the interaction energy $\langle\Psi_M^\pm|H_{AB}|\Psi_M^\pm\rangle = E_\pm - kE_B \equiv \varepsilon^\pm$. Therefore, from

$$\int \mathrm{d}\mu(\Omega)|\chi(\Omega)|^2\langle\phi(\Omega)|H_{\boldsymbol{\sigma}}^\pm(\Omega;M)|\phi(\Omega)\rangle = \langle\Psi_M^\pm|H_{AB}|\Psi_M^\pm\rangle \tag{22}$$

we define the effectively-local parametric Hamiltonians $H_{\boldsymbol{\sigma}}^\pm(\Omega;M)$ as

$$H_{\boldsymbol{\sigma}}^\pm(\Omega;M) = \varepsilon^\pm\frac{\boldsymbol{\sigma}}{2}\cdot\hat{\boldsymbol{n}}^\pm(\Omega;M)\ , \tag{23}$$

with $\hat{\boldsymbol{n}}^\pm(\Omega;M) = (\sin\Theta_M^\pm\cos\varphi, \sin\Theta_M^\pm\sin\varphi, \cos\Theta_M^\pm)$, $\Theta_M^\pm$ from Eq.(20), and $\varepsilon^\pm = \pm g(\tilde{S}\mp 1)/2N$. The above result is readily interpreted by saying that when the star is in one of its eigenstates, the qubit behaves as if it were a closed system under the effect of a magnetic field whose direction is distributed according to the environmental quantum probability $|\chi_M^\pm(\Omega)|^2$. This also provides a key to the reading of Fig.1, whose lower panel visualizes the qubit response to the application of a field with direction distributed as shown in the upper panel. Picturing the qubit in terms of the above "effective" magnetic field, one is naturally led to the last part of our work, namely the crossover from a quantum to a classical environment.

IV. THE LARGE-S LIMIT.

From a direct calculation one can see that the environmental probability distribution $p_M^\pm(\theta)$ is the properly normalized Q-representation of the environment density matrix and that

$$p_M^\pm(\theta) \xrightarrow{S\to\infty} \delta(\theta-\theta_M)\ ; \tag{24}$$

this means that when the environment is driven to its classical limit, the qubit parametrized states (19) collapse into the GS and excited state (according to whether the original state belonged to the GS "-" or excited "+" multiplet) of the effectively local Hamiltonian $H_{\boldsymbol{\sigma}}^{\rm cl} = \frac{g}{2}\boldsymbol{\sigma}\cdot\hat{\boldsymbol{n}}(\theta_M,\varphi)$, which is consistently recognized as the operator the two effectively local Hamiltonians $H_{\boldsymbol{\sigma}}^\pm(\Omega;M)$ collapse into, as $\theta\to\theta_M$. Notice that $S\to\infty$ implies $S/N\to 1/2$, so that $\varepsilon^\pm$ consistently converge to the physical coupling g. The local "qubit in a field" Hamiltonian is hence derived from the non-local model of an OQS, namely a qubit interacting with a magnetic environment via an isotropic quantum Heisenberg-like interaction, when the classical limit of the environment is taken. Despite such conclusion could be reached also by qualitative arguments based on mean-field and/or semiclassical treatments, the derivation here presented allows us to exactly obtain (rather than taking them as phenomenological parameters) the polar angle of the field direction and the intensity of the effective magnetic field, which indeed depend on the quantum number M of the state the global system is in, $|\Psi_M^\pm\rangle$, and the interaction coupling g, respectively.

The formal content of the above discussion allows us to take our last step forward and relate the entanglement of $|\Psi_M^\pm\rangle$ to the Berry's phase emerging in the corresponding effectively-local model. Let us first remind that the environmental coherent-state probability distribution, $p_M^\pm(\theta)$, does not depend on φ, which is therefore left undetermined by the choice of the eigenstate $|\Psi_M^\pm\rangle$. On the other hand, φ enters the expression of the qubit parametrized states (19), which means that an evolution of the environmental coherent states such that $\theta = \text{const}$ and φ varies, despite leaving the state of the global system unchanged, makes the qubit states evolve accordingly. Indeed, one such internal evolution corresponds, in the large-S limit where the effectively local Hamiltonian $H_{\boldsymbol{\sigma}}^{\rm cl}$ is defined, to the adiabatic precession of $\hat{\boldsymbol{n}}$ around the z axis, and give rise, when closed paths are considered, to a Berry's geometric phase that reads

$$\gamma_\pm = \mp\pi(1-\cos\theta_M)\ , \tag{25}$$

where the ± sign refers to the qubit being in the ground or excited state of $H_{\boldsymbol{\sigma}}^{\rm cl}$. On the other hand, the entanglement between the qubit and its environment when the star is in any of the states $|\Psi_M^\pm\rangle$ is, from Eq. (14),

$$\mathcal{E}_{\sigma S} = -h\left[\frac{\gamma_\pm}{2\pi}\right]\ , \tag{26}$$

i.e. the binary entropy of the 2π normalized Berry' phase. Notice that, being $h[x]=h[1-x]$, the dependence on ± is not transferred into the entanglement, as it must be the case, according to Eq.(14).

We understand the above result as follows: When we describe the system in terms of only one quantum system (the qubit), effectively reducing the environment to an external field which is treated at a classical level, we can no longer speak about entanglement (there cannot be entanglement if a system is not composite). However, the entangled structure of $|\Psi\rangle_M^\pm$ causes the dependence on φ to be conveyed from the environment to the qubit, so that a parametric dependence survives also in the large-S limit and makes the emergence of the Berry's phase possible. To this respect, notice that φ-paths in the parameters space that give rise to no Berry's phase ($\theta_M = 0,\pi$) consistently derive from *separable* states in the original fully-quantum mechanical description.

The result embodied by Eq.(26) pertains to the analysis of the relation between geometrical properties of quantum systems and the structure, or dynamical evolution, of their states [31, 32]. Of particular relevance, in the context of this work, are studies about the relation between Berry's adiabatic phases, and geometric projective phases [40], possibly arising in composite systems, and the entanglement relative to their partitions[41–44]; equally relevant are the non-Abelian generalizations of geometric phase [45] in the case of mixed states undergoing both unitary and non-unitary evolutions [46, 47].

In order to accurately collocate our result in the overall picture, let us further comment upon its general meaning.

For the behaviour of a physical system to be possibly characterized by the emergence of geometrical effects, some geometrical space to be explored must exist. This can happen in two distinct physical setups: *i)* An OQS is described by an effective approach, i.e. in terms of a local parametric Hamiltonian, in which case the geometrical space is that of the Hamiltonian parameters. *ii)* An isolated system is prepared in a pure state which is not an eigenstate of its Hamiltonian, and unitarily evolves accordingly, in which case the geometrical space is the Hilbert space of the system itself. Notice that this second situation implies a preparation procedure based on an interaction between the system and some environment, such that the system is in a pure state after the procedure has completed. From the above statements a somehow puzzling picture emerges in the case of an OQS which is part of an "all-along isolated" system, by this meaning an isolated system which is in an eigenstate of its Hamitonian. In fact, geometrical effects can characterize its behaviour as obtained by the effective approach, but indeed have no formal placement in its exact description, as there is no geometrical space to be explored. On the other hand, the entanglement properties that characterize the exact description, formally have no meaning when the effective approach is adopted, as there cannot be entanglement with a classical environment. Therefore, neither of the above schemes can validate a procedure for relating the entanglement between parts of an "all-along isolated" system, and the geometrical effects possibly arising in the behaviour of any of such parts. In the parametric representation, which provides an interpolating scheme between the two approaches, the above procedure becomes available in the form of a crossover from a quantum to a classical environment, and, in the case of the spin system here considered, a relation as strict as that embodied by Eq.(26) finally emerges.

V. CONCLUSIONS

From a general perspective, this work clarifies why modelling a closed quantum system with a parametric Hamiltonian implies the existence of an environment ("the rest of the Universe" to use Berry's words[30]) and shows that a non-trivial parametric dependence can arise if and only if such environment is entangled with the system itself. One of the most relevant consequence of the above statement is that the emergence of observable (i.e. gauge-invariant) quantities which are not eigenvalues of Hermitian operators of the system under analysis, such as the Berry's phase, turns out to be related not only to the fact that an environment exists [48], but specifically to the condition that the system be entangled with its environment. In fact, in the specific case of the spin-star here considered, Eq.(26) establishes that the entanglement between the qubit and its environment can be determined by measuring the observable Berry's phase characterizing the related model of a closed qubit in a magnetic field, which suggests a possible way for experimentally access entanglement properties via the observation of gauge-invariant phases.

We conclude by mentioning some possible developments of this work that we think are worth pursuing. First of all, one might exploit the peculiar properties characterizing the dynamics of generalized coherent states, in order to relate a possible dynamical evolution of the global system to that of the principal one. Indeed, the formal scheme here presented opens the possibility of using well-established approaches for dealing with quantum dynamics in phase space, such as the the path-integral formalism, the adiabatic perturbation theory, or the Born-Oppenheimer approximation, as tools for taking into account the effects of the environment on the principal system and vice versa [14, 49–52]. To this respect, notice that the approach here presented can be equivalently implemented in terms of generalized coherent states of any type, so that different physical systems can be taken into consideration[33], such as those belonging to the spin-bosons family. As for the spin-$\frac{1}{2}$ star with frustration, it is worth mentioning that different types of interaction between the environmental spins, in particular the antiferromagnetic Lieb-Mattis and Heisenberg-on-a-square-lattice ones, define exactly solvable models[53] that can be treated in the very same framework here proposed. This expands the set of real physical systems where to look for a possible experimental analysis of our results. Finally, we find particularly intriguing the idea of effectively studing the spin-$\frac{1}{2}$ star with frustration model by quantum simulators (see e.g. Ref.[54] and references therein, and Refs.[55, 56]): In fact, the possibility of tuning the interaction parameters, which is recognized as one of the key features of quantum simulators, might allow the variation of the value of S, by acting on the frustration ratio k/g, thus giving access to an experimental analysis of the crossover from a quantum to a classical environment.

Acknowledgments

The authors wish to thank T.J.G.Apollaro, L.Banchi, F.Bonechi, M.Tarlini, R.Vaia, Prof.A.Messina, and Prof.M.Long for fruitful discussions and valuable suggestions. Financial support of the Italian Ministry of Education, University, and Research in the framework of the 2008 PRIN program (Contract No. 2008PARRTS 003) is also acknowledged.

---

[1] H.P. Breuer and F. Petruccione. *The Theory of Open Quantum Systems.* Oxford University Press, USA, 2007.

[2] Á. Rivas and S.F. Huelga. *Open Quantum Systems: An Introduction.* SpringerBriefs in Physics. Springer, 2011.

[3] M.A. Nielsen and I.L. Chuang. *Quantum Computation and Quantum Information: 10th Anniversary Edition.* Cambridge University Press, 2000.

[4] P. Kaye, R. Laflamme, and M. Mosca. *An Introduction to Quantum Computing.* Oxford University Press, 2007.

[5] Geoffrey Hunter. Conditional probability amplitudes in wave mechanics. *International Journal of Quantum Chemistry*, 9(2):237–242, 1975.

[6] J Richter and A Voigt. The spin- 1/2 heisenberg star with frustration: numerical versus exact results. *Journal of Physics A: Mathematical and General*, 27(4):1139, 1994.

[7] Hong-Liang Deng and Xi-Ming Fang. Quantum phase transitions and quantum communication in a spin star system. *Journal of Physics B: Atomic, Molecular and Optical Physics*, 41(2):025503, 2008.

[8] N V Prokof'ev and P C E Stamp. Theory of the spin bath. *Reports on Progress in Physics*, 63(4):669, 2000.

[9] Heinz-Peter Breuer, Daniel Burgarth, and Francesco Petruccione. Non-markovian dynamics in a spin star system: Exact solution and approximation techniques. *Phys. Rev. B*, 70:045323, Jul 2004.

[10] A. Hutton and S. Bose. Mediated entanglement and correlations in a star network of interacting spins. *Phys. Rev. A*, 69:042312, Apr 2004.

[11] K. A. Al-Hassanieh, V. V. Dobrovitski, E. Dagotto, and B. N. Harmon. Numerical modeling of the central spin problem using the spin-coherent-state p representation. *Phys. Rev. Lett.*, 97:037204, Jul 2006.

[12] Francesca Palumbo, Anna Napoli, and Antonio Messina. Quantum correlations in a generalized spin star system. *Open Systems and Information Dynamics*, 13:309–314, 2006. 10.1007/s11080-006-9011-5.

[13] A Napoli, F Palumbo, and A Messina. Classical and nonclassical correlations in the dynamics of two spins coupled to a common spin bath. *Journal of Physics: Conference Series*, 36(1):154, 2006.

[14] Giuseppe Liberti, Rosa Letizia Zaffino, Franco Piperno, and Francesco Plastina. Entanglement of a qubit coupled to a resonator in the adiabatic regime. *Phys. Rev. A*, 73:032346, Mar 2006.

[15] Y. Hamdouni and F. Petruccione. Time evolution and decoherence of a spin-$\frac{1}{2}$ particle coupled to a spin bath in thermal equilibrium. *Phys. Rev. B*, 76:174306, Nov 2007.

[16] Hari Krovi, Ognyan Oreshkov, Mikhail Ryazanov, and Daniel A. Lidar. Non-markovian dynamics of a qubit coupled to an ising spin bath. *Phys. Rev. A*, 76:052117, Nov 2007.

[17] R. Hanson, V. V. Dobrovitski, A. E. Feiguin, O. Gywat, and D. D. Awschalom. Coherent dynamics of a single spin interacting with an adjustable spin bath. *Science*, 320(5874):352–355, 2008.

[18] E. Ferraro, H.-P. Breuer, A. Napoli, M. A. Jivulescu, and A. Messina. Non-markovian dynamics of a single electron spin coupled to a nuclear spin bath. *Phys. Rev. B*, 78:064309, Aug 2008.

[19] Michael Bortz, Sebastian Eggert, Christian Schneider, Robert Stübner, and Joachim Stolze. Dynamics and decoherence in the central spin model using exact methods. *Phys. Rev. B*, 82:161308, Oct 2010.

[20] Elena Ferraro. *Spin systems in fermionic and bosonic environments.* Diartimento di Scienze Fisiche ed Astronomiche, Università degli Studi di Palermo, 2010.

[21] Daniel Loss and David P. DiVincenzo. Quantum computation with quantum dots. *Phys. Rev. A*, 57:120–126, Jan 1998.

[22] B. E. Kane. A silicon-based nuclear spin quantum computer. *Nature*, 393:133, 1998.

[23] A. Imamoglu, D. D. Awschalom, G. Burkard, D. P. DiVincenzo, D. Loss, M. Sherwin, and A. Small. Quantum information processing using quantum dot spins and cavity qed. *Phys. Rev. Lett.*, 83:4204–4207, Nov 1999.

[24] John Schliemann, Alexander Khaetskii, and Daniel Loss. Electron spin dynamics in quantum dots and related nanostructures due to hyperfine interaction with nuclei. *Journal of Physics: Condensed Matter*, 15(50):R1809, 2003.

[25] W. A. Coish and Daniel Loss. Hyperfine interaction in a quantum dot: Non-markovian electron spin dynamics. *Phys. Rev. B*, 70:195340, Nov 2004.

[26] Davide Rossini, Tommaso Calarco, Vittorio Giovannetti, Simone Montangero, and Rosario Fazio. Decoherence induced by interacting quantum spin baths. *Phys. Rev. A*, 75:032333, Mar 2007.

[27] Łukasz Cywiński, Wayne M. Witzel, and S. Das Sarma. Pure quantum dephasing of a solid-state electron spin qubit in a large nuclear spin bath coupled by long-range hyperfine-mediated interactions. *Phys. Rev. B*, 79:245314, Jun 2009.

[28] P. Haikka, J. Goold, S. McEndoo, F. Plastina, and S. Maniscalco. Non-markovianity, loschmidt echo, and criticality: A unified picture. *Phys. Rev. A*, 85:060101, Jun 2012.

[29] Barry Simon. Holonomy, the quantum adiabatic theorem, and berry's phase. *Phys. Rev. Lett.*, 51:2167–2170, Dec 1983.

[30] M. V. Berry. Quantal phase factors accompanying adiabatic changes. *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*, 392(1802):45–57, 1984.

[31] Péter Lévay. The geometry of entanglement: metrics, connections and the geometric phase. *Journal of Physics A: Mathematical and General*, 37(5):1821, 2004.

[32] Dariusz Chruściński. Geometric aspects of quantum mechanics and quantum entanglement. *Journal of Physics: Conference Series*, 30(1):9, 2006.

[33] Wei-Min Zhang, Da Hsuan Feng, and Robert Gilmore. Coherent states: Theory and some applications. *Rev. Mod. Phys.*, 62:867–927, Oct 1990.

[34] Elliott H. Lieb. The classical limit of quantum spin systems. *Communications in Mathematical Physics*, 31:327–340, 1973. 10.1007/BF01646493.

[35] S. Boixo, L. Viola, and G. Ortiz. Generalized coherent states as preferred states of open quantum systems. *EPL (Europhysics Letters)*, 79(4):40003, 2007.

[36] Franklin Schroeck. *Quantum mechanics on phase space.* Fundamental theories of physics. Kluwer Academic, 1996.

[37] Gerardo Adesso and Fabrizio Illuminati. Entanglement

in continuous-variable systems: recent advances and current perspectives. *Journal of Physics A: Mathematical and Theoretical*, 40(28):7821, 2007.

[38] Ruggero Vasile, Paolo Giorda, Stefano Olivares, Matteo G. A. Paris, and Sabrina Maniscalco. Nonclassical correlations in non-markovian continuous-variable systems. *Phys. Rev. A*, 82:012313, Jul 2010.

[39] Elliott Lieb and Daniel Mattis. Ordering energy levels of interacting spin systems. *Journal of Mathematical Physics*, 3(4):749–751, 1962.

[40] Y. Aharonov and J. Anandan. Phase change during a cyclic quantum evolution. *Phys. Rev. Lett.*, 58:1593–1596, Apr 1987.

[41] Erik Sjöqvist. Geometric phase for entangled spin pairs. *Phys. Rev. A*, 62:022109, Jul 2000.

[42] D. M. Tong, E. Sjöqvist, L. C. Kwek, C. H. Oh, and M. Ericsson. Relation between geometric phases of entangled bipartite systems and their subsystems. *Phys. Rev. A*, 68:022106, Aug 2003.

[43] Reinhold A. Bertlmann, Katharina Durstberger, Yuji Hasegawa, and Beatrix C. Hiemayr. Berry phase in entangled systems: A proposed experiment with single neutrons. *Phys. Rev. A*, 69:032112, Mar 2004.

[44] B. Basu. Relation between concurrence and berry phase of an entangled state of two spin-$\frac{1}{2}$ particles. *Europhys. Lett.*, 73(6):833–838, 2006.

[45] Armin Uhlmann. Parallel transport and quantum holonomy along density operators. *Reports on Mathematical Physics*, 24(2):229 – 240, 1986.

[46] Erik Sjöqvist, Arun K. Pati, Artur Ekert, Jeeva S. Anandan, Marie Ericsson, Daniel K. L. Oi, and Vlatko Vedral. Geometric phases for mixed states in interferometry. *Phys. Rev. Lett.*, 85:2845–2849, Oct 2000.

[47] D. M. Tong, E. Sjöqvist, L. C. Kwek, and C. H. Oh. Kinematic approach to the mixed state geometric phase in nonunitary evolution. *Phys. Rev. Lett.*, 93:080405, Aug 2004.

[48] Raffaele Resta. Manifestations of berry's phase in molecules and condensed matter. *Journal of Physics: Condensed Matter*, 12(9):R107, 2000.

[49] Hiroshi Kuratsuji and Shinji Iida. Effective action for adiabatic process. *Progress of Theoretical Physics*, 74(3):439–445, 1985.

[50] Hiroshi Kuratsuji. Geometric canonical phase factors and path integrals. *Phys. Rev. Lett.*, 61:1687–1690, Oct 1988.

[51] S. Teufel. *Adiabatic Perturbation Theory in Quantum Dynamics*. Number No. 1821 in Lecture Notes in Mathematics. Springer, 2003.

[52] Gianluca Panati, Herbert Spohn, and Stefan Teufel. The time-dependent born-oppenheimer approximation. *ESAIM: Mathematical Modelling and Numerical Analysis*, 41(02):297–314, 2007.

[53] Johannes Richter, Andreas Voigt, Sven E Kroeger, and Claudius Gros. The spin- 1/2 heisenberg star with frustration: Ii. the influence of the embedding medium. *Journal of Physics A: Mathematical and General*, 29(4):825, 1996.

[54] Iulia Buluta and Franco Nori. Quantum simulators. *Science*, 326(5949):108–111, 2009.

[55] K. Kim, M.-S. Chang, S. Korenblit, R. Islam, E. E. Edwards, J. K. Freericks, G.-D. Lin, L.-M. Duan, and C. Monroe. Quantum simulation of frustrated ising spins with trapped ions. *Nature*, 465:590–593, 2010.

[56] Liang-Hui Du, Xingxiang Zhou, Yong-Jian Han, Guang-Can Guo, and Zheng-Wei Zhou. Strongly coupled josephson-junction array for simulation of frustrated one-dimensional spin models. *Phys. Rev. A*, 86:032302, Sep 2012.